

DETERMINATION OF THE MINIMAL LENGTH BY MICROSCOPIC BLACK HOLE TEMPERATURE

Vladan Panković^{*,#}

^{*}Department of Physics, Faculty of Sciences, 21000 Novi Sad,
Trg Dositeja Obradovića 4. , Serbia, vdpan@neobee.net

[#]Gimnazija, 22320 Indjija, Trg Slobode 2a, Serbia

PACS number: 04.70.Dy

Abstract

Generalizing results of our previous work (where classical kinetic energy has been used) in this work (where ultra-relativistic kinetic energy is used) we suggest an original variant of the determination of minimal length (corresponding to Plank length) by formation of a microscopic (tiny) black hole. Like to some previous authors we use Heisenberg coordinate-momentum uncertainty relation, on the one hand. But, instead of metric fluctuation (obtained by second derivative in Einstein equations) that generalizes uncertainty relation by an additional term, used by previous authors, we use Hawking temperature of the black hole and standard Heisenberg coordinate-momentum uncertainty relation.

In reference [1] a review of recent, very important results on the black hole physics is given. Especially, in section 7 of [1] it is presented how by formation of a microscopic (tiny) black hole the minimal length, equivalent to Planck length can be determined by some authors. Simply speaking, by measurement of the coordinate of a quantum system, or, simply speaking, a particle, standard, Heisenberg coordinate-momentum uncertainty relation must be satisfied. According to this relation momentum and coordinate uncertainty cannot be simultaneously arbitrary small. But, by arbitrary accurate measurement of coordinate corresponding coordinate uncertainty can be arbitrary small. For this reason standard Heisenberg coordinate-momentum uncertainty relation does not predict non-zero minimal length. However, coordinate measurement, roughly speaking, considers an interaction of given particle with a measuring particle at a corresponding distance (length). If given distance, by sufficiently accurate coordinate measurement, becomes equal or smaller than Schwarzschild radius for measured and measuring particle, then a microscopic (tiny) Schwarzschild black hole will appear. It, according to Einstein equations, will cause additional, nonstandard, metric fluctuation or uncertainty, whose value is proportional to product of the particle momentum and quadrate of Planck length. Or, there is a generalization of the coordinate-momentum uncertainty relation so that minimal length is equivalent to Plank length (smaller

length cannot be stably measurable since corresponding measuring procedure causes microscopic (tiny) black hole formation and non-removable metric fluctuations).

In this work, generalizing results of our previous work [2] (where classical kinetic energy has been used) we shall suggest (by use of more accurate, ultra-relativistic kinetic energy) an original variant of the determination of minimal length (corresponding to Plank length) by formation of a microscopic (tiny) black hole. Like to some previous authors we shall use Heisenberg coordinate-momentum uncertainty relation, on the one hand. But, instead of metric fluctuation (obtained by second derivative in Einstein equations) that generalizes uncertainty relation by an additional term, used by previous authors, we shall use Hawking temperature of the black hole and standard Heisenberg coordinate-momentum uncertainty relation.

We shall start from well-known, standard, Heisenberg momentum-coordinate uncertainty relation

$$\Delta p \Delta x \geq \frac{\hbar}{2} \quad (1)$$

where, for measured quantum system, i.e. particle, Δp represents the momentum uncertainty (or, standard deviation), Δx - coordinate uncertainty (or, standard deviation), and $\hbar$ - reduced Planck constant. As it has been mentioned previously, given uncertainty relation states that momentum uncertainty and coordinate uncertainty cannot be simultaneously arbitrary small. But given relation does not forbid such arbitrary accurate measurement of coordinate within which coordinate uncertainty can be arbitrary small. It implies possibility of infinity small length. Namely, by physically relevant measurement of an observable, e.g. coordinate, its average value cannot be smaller than its standard deviation. But, since in sufficiently accurate coordinate measurement coordinate uncertainty can be arbitrary small, arbitrary small coordinate, i.e. length can be physically relevant.

Uncertainty relation (1) implies

$$\Delta p^2 \simeq \langle p^2 \rangle - \langle p \rangle^2 \geq \frac{\hbar^2}{4\Delta x^2} \quad (2)$$

and, further,

$$\langle p^2 \rangle \geq \frac{\hbar^2}{4\Delta x^2}. \quad (3)$$

where $\langle \rangle$ denotes average value of corresponding observable.

For an ultra-relativistic quantum system, for which average value of the total energy $\langle E \rangle$ is practically equivalent to average value of the kinetic energy $\langle E_k \rangle$ or to $\langle p^2 \rangle^{\frac{1}{2}} c$ where c represents the speed of light, (3) simply implies

$$\Delta x \geq \frac{\hbar c}{2 \langle E_k \rangle} \quad (4)$$

Suppose that for sufficiently accurate measurement of the coordinate of particle, i.e. for sufficiently small Δx , given particle becomes completely a Schwarzschild black hole. Then average kinetic energy of the particle must be equivalent to thermal energy of the black hole. (Precisely speaking, since coordinate and momentum represent non-commutative observables then, according to standard quantum mechanical propositions, accurate measurement of coordinate causes significant statistical fluctuations of the momentum. Also, since, in ultra-relativistic case, momentum is practically proportional to kinetic and total energy, given accurate coordinate measurement causes significant statistical fluctuation of the kinetic energy too. For this reason it seems quite naturally

that given statistically averaged kinetic energy of the particle be considered as the thermal energy, i.e. energy of the statistical fluctuations of the black hole.) This thermal energy corresponding to chaotic oscillations at black hole horizon surface can be presented as $2(\frac{kT}{2}) = kT$ where 2 refers on two degree of freedom at horizon surface, while k represents Boltzmann constant and T - Hawking temperature. It yields

$$\Delta x \geq \frac{\hbar c}{2kT}. \quad (5)$$

As it is well-known, Hawking temperature for Schwarzschild black hole with mass m equals

$$T = \frac{\hbar c^3}{8k\pi Gm} \quad (6)$$

where G represents the gravitational constant. Then, introduction of (6) in (5) yields

$$\Delta x \geq \frac{4\pi Gm}{c^2} = \frac{4\pi Gmc}{c^3} = 4\pi \left(\frac{G\hbar}{c^3}\right) \frac{2\pi}{\lambda} = 8\pi^2 \left(\frac{G\hbar}{c^3}\right) \frac{1}{\lambda} = 8\pi^2 L_P^2 \frac{1}{\lambda} \quad (7)$$

where $\lambda = \frac{2\pi\hbar}{mc}$ represents corresponding de Broglie wave length while $L_P = \left(\frac{G\hbar}{c^3}\right)^{\frac{1}{2}}$ represents Planck length. Since, in given measurement procedure, Δx is practically equivalent to de Broglie wave length, (7) implies

$$\Delta x^2 \geq 8\pi^2 L_P^2 \quad (8)$$

and finally

$$\Delta x \geq 8^{\frac{1}{2}} \pi L_P \sim L_P. \quad (9)$$

In this way we did an original variant of the determination of minimal length (corresponding to Plank length) by formation of a microscopic (tiny) black hole. Like to some previous authors we use Heisenberg coordinate-momentum uncertainty relation, on the one hand. But, instead of metric fluctuation (obtained by second derivative in Einstein equations) that generalizes uncertainty relation by an additional term, used by previous authors, we used Hawking temperature of black hole and standard Heisenberg coordinate-momentum uncertainty relation.

References

- [1] S. Hossenfelder, *What black holes can teach us*, hep-ph/0412265 and references therein
- [2] V. Pankovic, *How we can do a microscopic (tiny) black hole and determine minimal length*, gr-qc/0901.3004